

Fold

Fold

Fold

Fold



FOR HEALTHCARE PROVIDERS

⚠️ This Patient has received ELREXFIO.

+ IMPORTANT SAFETY INFORMATION YOU SHOULD KNOW:
 ELREXFIO therapy can cause cytokine release syndrome (CRS) or neurologic toxicity including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) which may be fatal or life threatening. CRS may involve multiple organ systems.

FOR HEALTHCARE PROVIDERS

• First Treatment Dose _____

• Step-up Dose 2 _____

• Step-up Dose 1 _____

Dates of ELREXFIO Injections : _____

Healthcare Setting Name: _____

After Hours Phone Number: _____

Office Phone Number: _____

Name of ELREXFIO treating Oncologist: _____

ELREXFIO™ REMS
Patient Wallet Card
 1-844-923-7845 www.ELREXFIOREMS.com



Carry this card with you at all times.
SHOW THIS CARD to any healthcare provider involved
 in your care and if you go to the emergency room.



**IMPORTANT SAFETY INFORMATION FOR PATIENTS
 RECEIVING TREATMENT WITH ELREXFIO**

FOR THE PATIENT

Call your healthcare provider or get emergency help
 right away if you have any of these symptoms:

- | | |
|---|--|
| <ul style="list-style-type: none"> • Fever 100.4°F (38°C) or higher • Trouble breathing • Chills • Dizziness or light-headedness • Fast heartbeat • Headache • Agitation, trouble staying awake, confusion or disorientation, seeing or hearing things that are not real | <ul style="list-style-type: none"> • Trouble speaking, thinking, remembering things, paying attention, or understanding things • Problems walking or muscle weakness • Shaking (tremors), loss of balance, or muscle spasms • Numbness and tingling (feeling like "pins and needles") • Burning, throbbing, or stabbing pain • Changes in your handwriting |
|---|--|

IMPORTANT TO REMEMBER:



If you have **any** of these symptoms call your doctor or seek emergency medical attention right away! These are not all of the possible symptoms of ELREXFIO. Tell your doctor if you have any symptom that bothers you or does not go away.

You should be hospitalized after administration of each of the first 2 step-up doses within the step-up dosing schedule.

You should always ask your doctor about taking other medications while taking ELREXFIO.

Print out, cut along the dotted lines and fold in half and half again.