

+ IMPORTANT SAFETY INFORMATION FOR PATIENTS RECEIVING TREATMENT WITH TECVAYLI or TALVEY

Carry this card with you at all times. SHOW THIS CARD to any healthcare professional involved in your care and if you go to the emergency room.

TECVAYLI[®] and TALVEY[®] REMS PATIENT WALLET CARD

FOR HEALTHCARE PROFESSIONALS

+ IMPORTANT SAFETY INFORMATION YOU SHOULD KNOW: TECVAYLI and TALVEY therapy can cause cytokine release syndrome (CRS) or neurologic toxicity, including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) which may be fatal or life-threatening. CRS may involve multiple organ systems.

Complete the Patient Wallet Card based on the drug prescribed and provide the Patient Wallet Card to the patient.

⚠ This Patient has received: ☐ TECVAYLI ☐ TALVEY (Please check one.)

Name of Treating Oncologist: _____

After Hours Phone Number: _____ Office Phone Number: _____

Healthcare Setting Name: _____

Dosing Schedule: ☐ Weekly Dosing ☐ Bi-Weekly Dosing (Please check one.)

Dates of Injections: _____

• Step-up Dose 1: _____ • Step-up Dose 2: _____

• Step-up Dose 3*: _____ • First Treatment Dose: _____

*Applies only if patient is receiving TALVEY at a treatment dose of 0.8 mg/kg

FOR THE PATIENT

Call your healthcare professional or get emergency help right away if you recognize any of these symptoms:

Symptoms of Cytokine Release Syndrome (CRS):

- Fever (100.4°F or higher)
- Difficulty breathing
- Chills
- Fast heartbeat
- Dizziness or light-headedness
- Feeling anxious
- Confusion or restlessness
- Headache

Symptoms of neurologic problems:

- Headache
- Feeling confused or disoriented
- Being less alert or aware
- Feeling sleepy
- Low energy (lethargy)
- Slowness or difficulty thinking
- Seizures
- Memory Loss
- Jerking movements
- Rigid muscles
- Feeling restless
- Difficulty of coordination (affecting movement and balance)
- Numbness and tingling (feeling like “pins and needles”)
- Trouble speaking or understanding speech
- Muscle spasms
- Shaking (tremors)
- Problems walking
- Hearing Loss
- Muscle weakness in your body or face
- Double vision
- Burning, throbbing or stabbing pain
- Fast eye movements that cannot be controlled

You should always ask your doctor about taking other medications while taking TECVAYLI or TALVEY.



IMPORTANT TO REMEMBER: You may be asked to stay in the hospital for 48 hours after administration of all doses within the step-up dosing schedule. If you have any of these symptoms, call your doctor or seek emergency medical attention right away! These are not all of the possible symptoms of TECVAYLI or TALVEY. Tell your doctor if you have any symptom that bothers you or does not go away.